arXiv:1302.2069v1 [quant-ph] 8 Feb 2013

Macroscopic superpositions in Bose-Josephson junctions: controlling decoherence due to atom losses

K. Pawlowski,[1, 2, 3] D. Spehner,[4, 5, *] A. Minguzzi,[5] and G. Ferrini[6]

[1] *Center for Theoretical Physics PAN, Al. Lotników 32/46, 02-668 Warsaw, Poland*
[2] *5. Physikalisches Institut, Universität Stuttgart,*
Pfaffenwaldring 57, D-70569 Stuttgart, Germany
[3] *Laboratoire Kastler Brossel, Ecole Normale Supérieure, 45, rue d'Ulm, 75231 Paris, France*
[4] *Université Grenoble 1 and CNRS, Institut Fourier UMR5582, B.P. 74, 38402 Saint Martin d'Hères, France*
[5] *Université Grenoble 1 and CNRS, Laboratoire de Physique et Modélisation*
des Milieux Condensés UMR5493, B.P. 166, 38042 Grenoble, France
[6] *Laboratoire Kastler Brossel, Université Pierre et Marie Curie, 4 place de Jussieu, 75005 Paris, France*

(Dated: February 11, 2013)

We study how macroscopic superpositions of coherent states produced by the nondissipative dynamics of binary mixtures of ultracold atoms are affected by atom losses. We identify different decoherence scenarios for symmetric or asymmetric loss rates and interaction energies in the two modes. In the symmetric case the quantum coherence in the superposition is lost after a single loss event. By tuning appropriately the energies we show that the superposition can be protected, leading to quantum correlations useful for atom interferometry even after many loss events.

PACS numbers: 03.75Gg, 42.50.Lc, 03.75.Lm, 03.75.Mn, 03.65.Ud, 67.85.Hj, 03.75.Dg

In large systems, macroscopic superpositions of quantum states have extremely small decoherence times, making them impossible to observe [1]. The largest superpositions generated so far are superpositions of coherent states (CS) of a cavity field, and their progressive transformation into statistical mixtures as time evolves has been observed [2]. The first proposal to generate a macroscopic superposition of CS (MSCS) with light consisted in sending photons through a medium presenting a strong Kerr non-linearity [3]. In such media, the dynamical phases of Fock states are nonlinear in the photon number, thus the phase of an initial CS is split. In metastable vapors of ultracold bosonic atoms, interactions between atoms lead to similar nonlinearities, whose strength can be tuned experimentally by using Feshbach resonances [4, 5]. By trapping optically the condensed atoms in a double-well potential one realizes an external Bose-Josephson junction (BJJ); an internal BJJ is obtained by trapping in a single well atoms in two distinct hyperfine states coupled by a resonant field. In analogy with light, the nonlinear dynamics generates MSCSs after a sudden quench of the tunnel amplitude (for external BJJs) or a switch off of the coupling field (for internal BJJs) [6, 7]. Whereas MSCSs of light are presumably destroyed after a single photon loss [2] the situation is less clear for atoms. Up to now only squeezed states – which are produced at earlier times than MSCSs – have been observed in BJJs with a few hundreds of atoms [8–10]. In order to know if future experiments could produce MSCSs in BJJs, it is desirable to study how robust they are with respect to decoherence.

Decoherence effects due to atom losses and phase noise on squeezed states have been analyzed in detail [11–13]. For what concerns MSCSs, only decoherence caused by noise due to photon scattering [14] and phase noise [15] has been studied so far. Under current experimental conditions the first noise is negligible and the second one can be reduced by using spin-echo [10, 16]; in contrast, atom losses are unavoidable.

In this letter, we investigate how MSCSs generated in BJJs are affected by atom losses. Even though our analysis also applies to one- and three-body losses [17] we focus on two-body losses, due to scattering of two atoms in the magnetic trap which changes their spin and gives them enough kinetic energy to be ejected from the trap. The latter processes are particularly detrimental in the experiments of Refs. [9, 10]. We first analyze the dynamics of a lossy BJJ from the point of view of state conditioning, assuming that the total number $\hat{N}$ of condensed atoms can be measured precisely, both initially and at the MSCS formation time t_q. We study how much coherence is destroyed by a single loss event occurring at a random time between 0 and t_q. We find quite different answers depending on the degree of asymmetry between the loss rates and interactions energies in the two modes of the junction. Finally, we show that for strongly asymmetric losses one can protect the coherence of the MSCSs by suitably tuning the interaction energies, even after many loss events and in the absence of $\hat{N}$-measurement.

Quench dynamics in a BJJ in the presence of atom losses. We consider a two-mode BJJ in the quantum regime. Initially, N_0 atoms are in the ground state in the regime where tunneling dominates interactions, described by a spin coherent state

$$\begin{aligned}|\psi(0)\rangle &= |N_0;\theta,\phi\rangle \qquad (1)\\ &= \sum_{n_1=0}^{N_0}\begin{pmatrix} N_0\\ n_1\end{pmatrix}^{1/2}\frac{(e^{-\mathrm{i}\phi}\tan(\theta/2))^{n_1}}{[1+\tan^2(\theta/2)]^{N_0/2}}|n_1,N_0-n_1\rangle\end{aligned}$$

with $\theta = \pi/2$ and $\phi = 0$. Here $|n_1, n_2\rangle$ is the joint eigenstate of the number operators $\hat{n}_i$ in the mode $i = 1, 2$ (Fock state). The dynamics following a sudden quench of the tunnel energy to zero is given by the two-mode Bose-Hubbard Hamiltonian [18]

$$\hat{H}_0 = \sum_{i=1,2} \left(E_i \hat{n}_i + \frac{U_i}{2} \hat{n}_i (\hat{n}_i - 1) \right) + U_{12} \hat{n}_1 \hat{n}_2, \qquad (2)$$

where E_i and U_i are the internal energy (or, for external BJJs, the lowest energy level of the well i) and the interaction energy between two atoms in the mode i, respectively, and U_{12} is the inter-mode interaction energy (for external BJJs $U_{12} = 0$). Setting $\hat{n}_2 = N_0 - \hat{n}_1$, the interactions in Eq.(2) sum up to a non-linear term $\chi \hat{n}_1^2$, with $\chi = (U_1 + U_2 - 2U_{12})/2$. The time-evolved state at time $t_q = \pi/|\chi q|$ is a superposition of CSs, $|\psi^{(0)}(t_q)\rangle = e^{-\mathrm{i} t_q \hat{H}_0} |\psi(0)\rangle = \sum_{k=0}^{q-1} c_k |N_0; \frac{\pi}{2}, \phi_k\rangle$, with $\phi_{k+1} - \phi_k = 2\pi/q$ and $|c_k| = q^{-1/2}$ [3].

In the presence of two-body losses the markovian master equation for the density matrix $\hat{\rho}(t)$ of the condensed atoms reads [19, 20]

$$\frac{\mathrm{d}\hat{\rho}(t)}{\mathrm{d}t} = -\mathrm{i}\big[\hat{H}_0, \hat{\rho}\big] + \sum_{i=1,2} \gamma_i \Big(\hat{a}_i^2 \, \hat{\rho} \, (\hat{a}_i^2)^\dagger \qquad (3)$$

$$- \frac{1}{2} \big\{ \hat{n}_i (\hat{n}_i - 1), \hat{\rho} \big\} \Big) + \gamma_{12} \Big(\hat{a}_1 \hat{a}_2 \, \hat{\rho} \, \hat{a}_1^\dagger \hat{a}_2^\dagger - \frac{1}{2} \big\{ \hat{n}_1 \hat{n}_2, \hat{\rho} \big\} \Big)$$

where γ_i and γ_{12} are the loss rates of two atoms in the same mode i and of one atom in each mode, respectively, and $\{\cdot, \cdot\}$ denotes the anticommutator. Since Eq.(3) does not couple subspaces with different total atom numbers $\hat{N} = \hat{n}_1 + \hat{n}_2$ and $\hat{N} = N_0$ initially, at all times $\hat{\rho}(t)$ has the block structure $\hat{\rho}(t) = \sum_{N=0}^{N_0} w_N(t) \hat{\rho}_N(t)$, where $w_N(t)$ is the probability to have N atoms at time t and $\hat{\rho}_N(t)$ is the corresponding conditional state. This decomposition is naturally accounted for by quantum trajectories $t \mapsto |\psi_j(t)\rangle = |\widetilde{\psi}_j(t)\rangle / \|\widetilde{\psi}_j(t)\|$, $|\widetilde{\psi}_j(t)\rangle$ being the unnormalized wave function when j loss events occur at times $0 \leq s_1 \leq s_2 \leq \cdots \leq s_j \leq t$ in the channels $m_1, \cdots, m_j$ [2, 21, 22],

$$|\widetilde{\psi}_j(t)\rangle = e^{-\mathrm{i}(t-s_j)\hat{H}_{\mathrm{eff}}} \hat{M}_{m_j} e^{-\mathrm{i}(s_j - s_{j-1})\hat{H}_{\mathrm{eff}}} \hat{M}_{m_{j-1}} \cdots$$

$$\cdots e^{-\mathrm{i}\hat{H}_{\mathrm{eff}}(s_2 - s_1)} \hat{M}_{m_1} e^{-\mathrm{i}s_1 \hat{H}_{\mathrm{eff}}} |\psi(0)\rangle. \qquad (4)$$

For two-body losses one has three loss channels $m = 1$, 2, and 12, the corresponding jump operators are $\hat{M}_1 = \hat{a}_1^2$, $\hat{M}_2 = \hat{a}_2^2$, and $\hat{M}_{12} = \hat{a}_1 \hat{a}_2$. The dynamics between loss events is given by the effective Hamiltonian $\hat{H}_{\mathrm{eff}} = \hat{H}_0 - \mathrm{i}\hat{D}$; the damping operator $\hat{D} = (\sum_{i=1,2} \gamma_i \hat{n}_i (\hat{n}_i - 1) + \gamma_{12} \hat{n}_1 \hat{n}_2)/2$ describes the gain of information on the system resulting from the knowledge that no loss occurred. The conditional state $\hat{\rho}_N(t)$ after a detection of $N = N_0 - 2j$ atoms at time t is obtained by averaging over the j jump times s_k and channels m_k,

$$\widetilde{\rho}_N(t) \equiv w_N(t) \hat{\rho}_N(t) = \sum_{m_1, \cdots, m_j} \int_{0 \leq s_1 \leq \cdots \leq s_j \leq t} \mathrm{d}s_1 \ldots \mathrm{d}s_j$$

$$p^{(t)}_{m_1, \ldots, m_j}(s_1, \ldots, s_j; j) |\psi_j(t)\rangle\langle\psi_j(t)| \qquad (5)$$

where $p^{(t)}_{m_1, \ldots, m_j}(s_1, \ldots, s_j; j) = \gamma_{m_1} \ldots \gamma_{m_j} \|\widetilde{\psi}_j(t)\|^2$ is the joint distribution of the s_k, m_k, and j [22]. By further summing over the number of jumps j one gets the total density matrix $\hat{\rho}(t) = \sum_{N=0}^{N_0} w_N(t) \hat{\rho}_N(t)$ which is a solution of Eq.(3). In order to understand the effect of losses we analyze separately each N-atom sectors.

Density matrix in the subspace with N_0 atoms. When no loss occurs in the time interval $[0, t]$, from Eq.(4) we obtain the unnormalized conditional state $\widetilde{\rho}_{N_0}^{(\text{no loss})}(t) = |\widetilde{\psi}_0(t)\rangle\langle\widetilde{\psi}_0(t)|$ in the Fock basis,

$$\langle n_1, n_2 | \widetilde{\rho}_{N_0}^{(\text{no loss})}(t) | n_1', n_2' \rangle$$

$$= e^{-t[d_{N_0}(n_1) + d_{N_0}(n_1')]} \langle n_1, n_2 | \hat{\rho}^{(0)}(t) | n_1', n_2' \rangle, \quad (6)$$

where $d_{N_0}(n_1) = \frac{1}{2}(\gamma_1 + \gamma_2 - \gamma_{12})(n_1 - \overline{n}_1)^2$ up to an irrelevant constant, $2\overline{n}_1 = [\gamma_1 - \gamma_2 + N_0(2\gamma_2 - \gamma_{12})]/(\gamma_1 + \gamma_2 - \gamma_{12})$, and $\hat{\rho}^{(0)}(t) = |\psi^{(0)}(t)\rangle\langle\psi^{(0)}(t)|$ is the lossless density matrix. For $N_0 \gg 1$ the matrix elements of $\hat{\rho}^{(0)}(t)$ have Gaussian moduli peaked at $(n_1, n_1') = (N_0/2, N_0/2)$ with a width $\sim \sqrt{N_0}$. For symmetric loss rates $\gamma_1 = \gamma_2$ and $\gamma_{12} = 0$, this peak coincides with the center of the Gaussian damping factor in Eq.(6). Thus the MSCS formed at time t_q is affected by damping when $\gamma_1 \gtrsim \chi q / N_0$. In contrast, for $\gamma_2 = \gamma_{12} = 0$ the damping factor is centered at $(n_1, n_1') = (1/2, 1/2)$ and its effect on the MSCS sets in at the much smaller rate $\gamma_1 \approx \chi q / N_0^2$.

Density matrix in the subspace with $(N_0 - 2)$ atoms. We focus now on trajectories $t \mapsto |\psi_1(t)\rangle$ having one loss event in channel m at the random time $s \in [0, t]$. The instantaneous jump transforms a CS into a CS, $\hat{M}_m |N_0; \theta, \phi\rangle \propto |N_0 - 2; \theta, \phi\rangle$. This CS is rotated on the Bloch sphere by the evolution under the nonlinear effective Hamiltonian $\hat{H}_{\mathrm{eff}}$ due to changes in energy and damping after the jump, yielding

$$|\psi_1(t)\rangle \propto e^{-\mathrm{i}t\hat{H}_{\mathrm{eff}}} |N_0 - 2; \theta_m(s), \phi_m(s)\rangle \qquad (7)$$

where the random angles $\theta_m(s) = 2\arctan(e^{-s\delta_m})$ and $\phi_m(s) = 2s\chi_m$ depend on the random loss time s, the interaction energies $\chi_1 = U_1 - U_{12}$, $\chi_2 = -(U_2 - U_{12})$, $\chi_{12} = (U_1 - U_2)/2$, and the loss rate differences $\delta_1 = 2\gamma_1 - \gamma_{12}$, $\delta_2 = -(2\gamma_2 - \gamma_{12})$, and $\delta_{12} = \gamma_1 - \gamma_2$. Hence, apart from reducing N and producing damping, atom losses are identical to external θ and ϕ-noises rotating the state around the z-axis by a complex angle $\phi_m + \mathrm{i}\ln\tan(\frac{\theta_m}{2})$ [23]. These noises have fluctuations $\delta\theta_m \simeq |\delta_m| \min\{t, \delta s_m\}$ and $\delta\phi_m = 2|\chi_m| \min\{t, \delta s_m\}$, where δs_m is the loss-time fluctuation.

Tuning the energies at weak losses. Let us concentrate on the MSCS formation time $t = t_q$ and take $\gamma_{12} = 0$. We first consider *weak losses* $\gamma_m \lesssim q\chi/N_0$ and *symmetric energies* $U_1 = U_2$. In this regime the θ-noise is negligible since $\delta\theta_m$ is much smaller than the quantum fluctuations $\sim 1/\sqrt{N_0}$ of a CS. In contrast, one has large ϕ-fluctuations $\delta\phi_m = 2\pi/q$, equal to the inter-component phase separation of the MSCS (we use here



FIG. 1. (Color online) Modulus of the density matrix (8) in the subspace with $N_0 - 2$ atoms in the Fock basis at time t_2, obtained by an exact diagonalization of Eq.(3). Upper panels: symmetric losses rates ($\gamma_1 = \gamma_2 = \chi/(200\pi)$); lower panels: asymmetric losses ($\gamma_1 = 4\chi/(300\pi)$, $\gamma_2 = 0$). Left column: symmetric energies ($\chi_1 = -\chi_2 = \chi$); right column: asymmetric energies ($\chi_1 = 2\chi$, $\chi_2 = 0$). Other parameters: $\gamma_{12} = E_i = 0$, $N_0 = 100$.

$\chi_1 = -\chi_2 = \chi$ and $\delta s_m > t_q$). Such a strong ϕ-noise has a relatively small effect on the coherences, albeit it transforms the corresponding statistical mixture of CSs into a mixture of Fock states (phase relaxation) [15]. Actually, if $\gamma_2 = 0$ the off-diagonal elements of the conditional density matrix $\langle n_1, n_2 | \hat{\rho}_{N_0-2}(t_q) | n_1', n_2' \rangle$ are weakly affected by losses for $n_1' \neq n_1$ modulo 2, as shown in the appendix and in Fig. 1 (lower left panel). The global shift to values $n_1, n_1' < N_0/2$ is due to the Gaussian damping caused by $\hat{H}_{\rm eff}$ in Eq.(7). For $\gamma_1 = \gamma_2$, instead, $\hat{\rho}_{N_0-2}(t_2)$ is almost diagonal in the Fock basis due to a cancellation (occurring only for $q = 2$) when summing the contributions of the two channels $m = 1, 2$ (upper left panel in Fig.1). We now consider *asymmetric energies* $U_1 \neq U_2$. In order to keep $t_q = \pi/|\chi q|$ constant, we vary χ_1, χ_2 while fixing $2\chi = \chi_1 - \chi_2$. Interestingly, it is possible to protect against ϕ-noise one channel, say $m = 1$, by choosing $\chi_1 = 0$ and $\chi_2 = -2\chi$, to the expense of enlarging noise in the other channel. Then $\delta\phi_1 = 0$ and $\delta\phi_2 = 4\pi/q$. If only the first channel loses atoms, the conditional state after a single loss event is thus close to a MSCS with $N_0 - 2$ atoms, apart from the damping described in Eq.(6). For symmetric losses, $\hat{\rho}_{N_0-2}(t_q)$ has also large off-diagonal elements in the Fock basis due to probability one half to loose atoms in the protected channel (Fig. 1, right panels).

The ϕ-noise decreases at *intermediate loss rates* since the loss-time fluctuations δs_m decrease when increasing γ_m. Indeed, we find $\delta s_i \approx (2\gamma_i + \gamma_{12})^{-1} N_0^{-1}$ $(i = 1, 2)$ for $N_0 \gg 1$. Physically, at increasing γ_m the loss has more chance to occur at small times, while for small γ_m, s is equally distributed in $[0, t]$.

Subspaces with $N_0 - 2j$ atoms. The wavefunction $|\psi_j(t)\rangle$ after $j > 1$ jumps is still given by Eq.(7) upon replacing $N_0 - 2$ by $N_0 - 2j$ and the angles of the CS by $\theta^{(j)}$ and $\phi^{(j)} = \sum_{k=1}^{j} \phi_{m_k}(s_k)$, with $\tan(\frac{\theta^{(j)}}{2}) = \prod_{k=1}^{j} \tan(\frac{\theta_{m_k}(s_k)}{2})$. Thus the aforementioned effects persist; for weak symmetric losses, though, keeping coherence by switching off phase noise in one channel is harder since the probability that all jumps occur in that channel decreases exponentially with j. This means that for many loss events our proposal for protecting MSCSs is only efficient for strongly asymmetric rates.

Total quantum correlations and sub-shot noise atom interferometry. We now show that for completely asymmetric losses not only the conditional states but also the quantum correlations in the full density matrix $\hat{\rho}(t_q)$ can be protected by tuning the interactions. We measure the amount of quantum correlations with the quantum Fisher information $F(\hat{\rho})$, related to the best achievable phase precision in a Mach-Zehnder interferometer using $\hat{\rho}$ as input state by $(\Delta\varphi)_{\rm best} = 1/\sqrt{F(\hat{\rho})}$ [24]. Hence $F(\hat{\rho}) > \langle \hat{N} \rangle$ implies phase accuracy beyond the shot noise limit $(\Delta\varphi)_{\rm SN} = \langle \hat{N} \rangle^{-1/2}$ [25]. In our case $F_{\rm tot}(t) \equiv F[\hat{\rho}(t)] = \sum_{N=0}^{N_0} w_N(t) F_N(t)$ where $F_N(t)$ is the Fisher information of the conditional state $\hat{\rho}_N(t)$ in the subspace with N atoms [26]. We optimize $F_{\rm tot}(t)$ over all directions of the interferometer. In a lossless BJJ, the two-component superposition has the highest possible value $F_{\rm tot}(t_2) = N_0^2$, nearly twice larger than that of MSCSs with $q > 2$ components [27, 28].

Fig. 2 shows the Fisher information $F_N(t_2)$ in the $N = N_0 - 2$ subspace, corresponding to the state conditioned to a single loss event. For symmetric energies, the low values of F_N at small symmetric loss rates are direct consequences of the cancellation among channels. For $\gamma_2 = 0$ a much larger value is found, showing that the aforementioned non-vanishing inter-component coherences carry useful quantum correlations. At intermediate rates, $F_N(t_2)$ increases and reaches a maximum as a result of the reduced phase noise when $\gamma_1 = \gamma_2$ [30]. This reduction is clearly seen on the Husimi distributions $Q_N(\theta, \phi) = \frac{1}{\pi} \langle N; \theta, \phi | \hat{\rho}_N(t_2) | N; \theta, \phi \rangle$ which display a flat profile for small losses and two emerging peaks for larger losses (insets of Fig. 2). For asymmetric losses $\gamma_2 = 0$, by choosing $\chi_1 = 0$ we have $F_N(t_2) \to N^2$ in the small loss limit and the Husimi function has two peaks at the CS phases $\phi = \pm\pi/2$, in agreement with our prediction that the conditional state converges to a two-component superposition. At intermediate rates $F_N(t_2)$ decreases with γ_1 because the Gaussian damping compensates phase noise reduction. At large losses this damping transforms $\hat{\rho}_{N_0-2}(t_2)$ into a superposition of



FIG. 2. (Color online) Fisher information in the subspace with $N_0 - 2$ atoms as a function of the loss rate γ_1 (in semi-log scale) at time $t_2 = T/4$, with $T = 2\pi/\chi$. Solid black line: $\gamma_1 = \gamma_2$, $\chi_1 = -\chi_2 = \chi$; dotted red line: $\gamma_2 = 0$, $\chi_1 = -\chi_2 = \chi$; dashed blue line: $\gamma_2 = \chi_1 = 0$, $\chi_2 = -2\chi$; dot-dashed brown line: $\gamma_1 = \gamma_2$, $\chi_1 = 0$, $\chi_2 = -2\chi$. The optimization over the interferometer directions is done independently in the $(N_0 - 2)$-subspace. Insets: Husimi functions for the values of γ_i and χ_i indicated on the curves by black circles. Other parameters as in Fig. 1.



FIG. 3. (Color online) Total quantum Fisher information $F_{\text{tot}}(t)$ vs time t (in units of T). From top to bottom: $\gamma_1 T = 8/300$, $\gamma_2 = \chi_1 = 0$, $\chi_2 = -2\chi$ (blue dashed line); $\gamma_1 T = 8/300$, $\gamma_2 = 0$, $\chi_1 = -\chi_2 = \chi$ (red dotted line); $\gamma_1 T = \gamma_2 T = 1/100$, $\chi_1 = 0$, $\chi_2 = -2\chi$ (brown dot-dashed line); $\gamma_1 T = \gamma_2 T = 1/100$, $\chi_1 = -\chi_2 = \chi$ (black solid line). Other parameters as in Fig.1. Dashed horizontal line: F_{tot} for the highest squeezed state in the lossless case. Inset: average number of atoms vs. time for the same parameters.

Fock states with $n_1 = 0$ or 1 atoms in the first mode [17]. For symmetric losses and even N_0 , $\hat{\rho}_{N_0-2}(t_2)$ is transformed instead into the Fock state $|\frac{N}{2}, \frac{N}{2}\rangle$ having a larger Fisher information $\sim N^2/2$.

Our main result is presented in Fig. 3, which displays the total Fisher information $F_{\text{tot}}(t)$ obtained from an exact diagonalization of Eq.(3) for various rates and energies, keeping the same value of χ and of the mean number of atoms at time t_2 (see the inset). We find that for $\gamma_2 = 0$, tuning the energies has a strong effect on $F_{\text{tot}}(t)$ at times $\chi t \gtrsim N_0^{-1/2}$, yielding to larger Fisher informations than for squeezed states. Especially for the two-component superposition, quantum correlations are then preserved even after the loss of 20% of atoms.

Conclusions. We have shown that the interplay of atom losses and interactions in BJJs leads to different decoherence scenarios even after a single loss event. In particular, for strongly asymmetric two-body losses one can protect superpositions by tuning the interaction energies, leading to useful states for high-precision atom interferometry applications. Such asymmetric losses occur in internal BJJs with ^{87}Rb atoms in the hyperfine states $|F, m_F\rangle = |2, -1\rangle$ and $|1, 1\rangle$, subject respectively to fast two-body and slow one-body losses [16]. Finally, we note that ϕ-noise associated to the loss times s_k could be used to obtain an indirect measurement of s_k from the interferometric estimation of the phase rotation of MSCSs, yielding to an observation of quantum jumps like in cavity QED [2].

Appendix. Using Eq.(4) we find the distribution $p_m^{(t)}(s;1) = \gamma_m e^{-sG_m}[\cosh(s\delta_m)]^{N_0-2}\|e^{-it\hat{H}_{\text{eff}}}|N_0 - 2; \theta_m(s), 0\rangle\|^2 N_0(N_0-1)/4$ with $G_i = (2\gamma_i + \gamma_{12})N_0 - 2\gamma_i - 2\gamma_{12}$ and $G_{12} = (\gamma_1 + \gamma_2 + \gamma_{12})N_0 - 2\gamma_1 - 2\gamma_2 - \gamma_{12}$. Averaging over trajectories as in Eq.(5) yields

$$\langle n_1, n_2|\hat{\rho}_{N_0-2}(t)|n_1', n_2'\rangle \propto \sum_m \gamma_m C_m(t; n_1, n_1')\langle n_1, n_2|\widetilde{\rho}_{N_0-2}^{(\text{no loss})}(t)|n_1', n_2'\rangle, \quad (8)$$

where $\widetilde{\rho}_{N_0-2}^{(\text{no loss})}(t) = e^{-\mathrm{i}t\hat{H}_{\text{eff}}}|N_0{-}2; \frac{\pi}{2}, 0\rangle\langle N_0{-}2; \frac{\pi}{2}, 0|e^{\mathrm{i}t\hat{H}_{\text{eff}}^{\dagger}}$ is the conditional state having no loss in $[0, t]$ for an initial CS with $N = N_0 - 2$ atoms, and

$$C_m(t; n, n') = \frac{1 - e^{-t[G_m + \delta_m(n+n'-N_0+2) + 2\mathrm{i}\chi_m(n-n')]}}{G_m + \delta_m(n + n' - N_0 + 2) + 2\mathrm{i}\chi_m(n - n')}.$$

Eq.(8) shows that the density matrix conditioned to a single loss event is a MSCS with $N_0 - 2$ atoms modulated by the envelope $\sum \gamma_m C_m(t; n, n')$ and the damping factor of Eq.(6). For weak losses $\gamma_m \ll q\chi/N_0$ and $U_1 = U_2$, $|C_m(t_q; n, n')| \simeq \frac{\pi}{\chi q}|\operatorname{sinc}(\frac{\pi(n-n')}{q})|$ vanishes at the values $n' = n + pq$ ($p = \pm 1, \pm 2, \cdots$) corresponding to the nonzero off-diagonal elements of the statistical mixture of CSs $\sum_k |c_k|^2 |N; \frac{\pi}{2}, \phi_k\rangle\langle N; \frac{\pi}{2}, \phi_k|$, and decays like $|n' - n|^{-1}$ for the other off-diagonal elements encoding the coherences among the CSs [29]. Hence the coherences are non vanishing, except for $\gamma_1 = \gamma_2$ and $q = 2$, because then the contributions of the two channels cancel each other, $\sum_m \gamma_m C_m(t_2; n, n') \simeq 0$ for $n \neq n'$. Taking now $\chi_1 = 0$ and $\chi_2 = -2\chi$, one infers from $\sum_m \gamma_m C_m(t_2; n, n') \simeq \frac{\pi}{2\chi}(\gamma_1 + \gamma_2 \delta_{nn'})$ that $\hat{\rho}_{N_0-2}(t_2) \propto \widetilde{\rho}_{N_0-2}^{(\text{no loss})}(t_2)$ when $\gamma_2 = 0$. Therefore, the conditional state is a MSCS slightly modified by Gaussian damping. The above calculation can be generalized to trajectories with $j > 1$ jumps between 0 and t. For $\gamma_m t \ll 1$ and $j \ll N_0$, $\hat{\rho}_{N_0-2j}(t)$ is still given by Eq.(8) upon replacing $\sum \gamma_m C_m(t; n_1, n_1')$ by $[\sum_m \gamma_m C_m(t; n_1, n_1')]^j$.

Acknowledgements. We are grateful to K. Rzążewski, F. Hekking, P. Treutlein, C. Gross, and M. Oberthaler for helpful discussions. D.S and K.P acknowledges support from the project ANR-09-BLAN-0098-01, A.M. from the ERC Handy-Q project, and K.P. from the Polish Government Funds for the years 2010-2012 and the project "Decoherence in long range interacting quantum systems and devices" sponsored by the Baden-Württemberg Stiftung.

---

* Dominique.Spehner@ujf-grenoble.fr

[1] D. Giulini *et al.*, *Decoherence and the Appearance of a Classical Word in Quantum Theory* (Springer, Berlin, Heidelberg, 1996)

[2] S. Haroche and J.-M. Raimond, *Exploring the Quantum: Atoms, Cavities, and Photons* (Oxford University Press, Oxford, 2006)

[3] B. Yurke and D. Stoler, Phys. Rev. Lett. **57**, 13 (1986)

[4] U. Fano, Phys. Rev. **124**, 1866 (1961)

[5] H. Feshbach, Annals of Physics **5**, 357 (1958)

[6] F. Piazza, L. Pezzé, and A. Smerzi, Phys. Rev. A **78**, 051601 (2008)

[7] G. Ferrini, A. Minguzzi, and F. Hekking, Phys. Rev. A **78**, 023606 (2008)

[8] J. Estève, C. Gross, A. Weller, S. Giovanazzi, and M. K. Oberthaler, Nature **455**, 1216 (2008)

[9] F. Riedel, P. Böhi, Y. Li, T. W. Hänsch, A. Sinatra, and P. Treutlein, Nature **464**, 1170 (2010)

[10] C. Gross, T. Zibold, E. Nicklas, J. Estève, and M. K. Oberthaler, Nature **464**, 1165 (2010)

[11] A. Sinatra and Y. Castin, Eur. Phys. J. D **4**, 247 (1998); Y. Li, Y. Castin, and A. Sinatra, Phys. Rev. Lett. **100**, 210401 (2008); Y. Li, P. Treutlein, J. Reichel, and A. Sinatra, Eur. Phys. J. B **68**, 365 (2009)

[12] A. Sinatra, J.-C. Dornstetter, and Y. Castin, Front. Phys. **7**, 86 (2012)

[13] K. Pawłowski and K. Rzążewski, Phys. Rev. A **81**, 013620 (2010)

[14] Y. P. Huang and M. G. Moore, Phys. Rev. A **73**, 023606 (2006)

[15] G. Ferrini, D. Spehner, A. Minguzzi, and F. Hekking, Phys. Rev. A **82**, 033621 (2010)

[16] C. Gross, J. Phys. B **45**, 103001 (2012)

[17] K. Pawlowski, D. Spehner, A. Minguzzi, and G. Ferrini, *in preparation*

[18] G. J. Milburn, J. Corney, E. M. Wright, and D. F. Walls, Phys. Rev. A **55**, 4318 (1997)

[19] J. Anglin, Phys. Rev. Lett. **79**, 6 (1997)

[20] M. W. Jack, Phys. Rev. Lett. **89**, 140402 (2002); M. W. Jack, Phys. Rev. A **67**, 043612 (2003)

[21] H. Carmichael, *An Open System Approach to Quantum Optics* (Springer-Verlag, New York, 1991)

[22] J. Dalibard, Y. Castin, and K. Mølmer, Phys. Rev. Lett. **68**, 580 (1992)

[23] An analogous result has been found in [12] for weak losses, i.e., without θ-noise

[24] S. L. Braunstein and C. M. Caves, Phys. Rev. Lett. **72**, 3439 (1994)

[25] P. Hyllus, L. Pezzé, and A. Smerzi, Phys. Rev. Lett. **105**, 120501 (2010)

[26] For a given direction $\mathbf{n}$ of the interferometer, F_N is defined as $F_N = 2\sum_{k,l} \frac{(p_k-p_l)^2}{p_k+p_l} |\langle\varphi_k|\hat{J}_{\mathbf{n}}|\varphi_l\rangle|^2$ with $\hat{\rho}_N|\varphi_l\rangle = p_l|\varphi_l\rangle$, $\hat{J}_{\mathbf{n}}$ being the angular momentum operator in the direction $\mathbf{n}$.

[27] L. Pezzé and A. Smerzi, Phys. Rev. Lett. **102**, 100401 (2009)

[28] G. Ferrini, D. Spehner, A. Minguzzi, and F. Hekking, Phys. Rev. A **84**, 043628 (2011)

[29] More precisely, one can show [15] that the lossless density matrix $\langle n_1, n_2|\hat{\rho}^{(0)}(t_q)|n_1', n_2'\rangle$ in the Fock basis is the sum of two matrices, the first one having zero matrix elements for $n_1' \neq n_1$ modulo q and corresponding to the statistical mixture of CSs, and the second one having zero matrix elements for $n_1' = n_1$ modulo q and corresponding to the coherences $\sum_{k\neq k'} c_k c_{k'}^* |N; \frac{\pi}{2}, \phi_k\rangle\langle N; \frac{\pi}{2}, \phi_{k'}|$. The latter do almost not contribute to the Husimi function but are responsible for the high values of the Fisher information.

[30] Note, however, that the total Fisher information $F_{\mathrm{tot}}(t_q)$ decreases by increasing γ_m due to the rapid decay of the probability $w_{N_0-2}(t_q)$ of losing only two atoms.